



Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Folicur® 3.6 F Foliar Fungicide

For Use on Barley to suppress *Fusarium* head blight (scab).

SECTION 18 SPECIFIC EXEMPTION

File Symbol: 06-MT-01

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF MONTANA UNDER SECTION 18 SPECIFIC EXEMPTION

This labeling must be in possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire Directions for Use and Disclaimer of Warranties before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for Folicur® 3.6 F Foliar Fungicide (EPA Registration Number 264-752).

Any adverse effects resulting from the use of Folicur® 3.6 F Foliar Fungicide under this emergency exemption must be immediately reported to the State of Montana Department of Agriculture.

EFFECTIVE PERIOD:

This labeling becomes effective March 15, 2006 and expires on July 20, 2006.

DIRECTIONS FOR USE

A maximum of 45,500 acres of barley may be treated under this exemption.

RECOMMENDED APPLICATIONS

CROP	DISEASES	RATE OF FOLICUR® 3.6 F
Barley	<i>Fusarium</i> head blight (scab)	4 fl oz per acre

Apply Folicur® 3.6 F as a preventive foliar spray. For optimum disease suppression, the lowest rate of spray surfactant should be tank-mixed with Folicur® 3.6 F may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 3 gallons of spray solution per acre by aircraft spray equipment.

Application through any type of irrigation equipment is prohibited.

A maximum of 1 application per season may be made. A maximum of 4 fl oz of Folicur® 3.6 F may be applied per acre per crop season. Folicur® may be applied up through full head emergence (Feekes 10.5). Application may not be made within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Do not enter treated field for 12 hours after application.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater, estuarine and marine fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937)